Patient Information Sheet BUSPIRONE HYDROCHLORIDE, USP

Rx Only

Dispense with Patient Information Sheet available at:

https://www.oxford-rx.com/med-guides

HOW TO USE: Buspirone Hydrochloride Tablets, USP 15 mg and 30 mg Tablets


in convenient adjustable dosage tablet form

Response to buspirone varies among individuals. Your physician may find it necessary to adjust your dosage to obtain the proper response.

This adjustable dosage tablet design makes dosage adjustments easy. Each tablet is scored and can be broken accurately to provide any of the following dosages.

If your doctor prescribed the 30-mg tablet:	If your doctor prescribed the 15-mg tablet:
10 10 10 30 mg (the entire tablet)	0 9 3 15 mg (the entire tablet)
20 mg (two thirds of a tablet)	10 mg (two thirds of a tablet)
10 mg (one third of a tablet)	0 5 mg (one third of a tablet)
10 1 15 mg (one half of a tablet)	7.5 mg (one half of a tablet)

To break a adjustable dosage tablet accurately and easily, hold the tablet between your thumbs and index fingers close to the appropriate tablet score (groove) as shown in the photo. Then, with the tablet score facing you, apply pressure and snap the tablet segments apart (segments breaking incorrectly should not be used).



Manufactured by: Oxford Pharmaceuticals LLC 301 Leaf Lake Parkway, Birmingham, AL 35211 USA