

**Patient Information Sheet**  
**BUSPIRONE HYDROCHLORIDE, USP**

**Rx Only**


Dispense with Patient Information Sheet available at:  
<https://www.oxford-rx.com/med-guides>

**HOW TO USE:**  
**Buspirone Hydrochloride Tablets, USP**  
**15 mg and 30 mg Tablets**  
in convenient adjustable dosage tablet form


Response to buspirone varies among individuals. Your physician may find it necessary to adjust your dosage to obtain the proper response.

This adjustable dosage tablet design makes dosage adjustments easy. Each tablet is scored and can be broken accurately to provide any of the following dosages.

If your doctor  
prescribed  
the 30-mg  
tablet:



30 mg  
(the entire  
tablet)



20 mg  
(two thirds  
of a tablet)




10 mg  
(one third  
of a tablet)



15 mg  
(one half  
of a tablet)

If your doctor  
prescribed  
the 15-mg  
tablet:



15 mg  
(the entire  
tablet)



10 mg  
(two thirds  
of a tablet)




5 mg  
(one third  
of a tablet)



7.5 mg  
(one half  
of a tablet)

To break an adjustable dosage tablet accurately and easily, hold the tablet between your thumbs and index fingers close to the appropriate tablet score (groove) as shown in the photo. Then, with the tablet score facing you, apply pressure and snap the tablet segments apart (segments breaking incorrectly should not be used).



Manufactured by: Oxford Pharmaceuticals LLC  
301 Leaf Lake Parkway, Birmingham, AL 35211 USA